



## Illinois Wesleyan University Digital Commons @ IWU

---

John Wesley Powell Student Research  
Conference

1997, 8th Annual JWP Conference

---

Apr 12th, 10:45 AM - 11:00 AM

# Adapting Daubechies' Wavelet Iteration Scheme to Biorthogonal Scaling Functions

Nathan Mueggenburg  
*Illinois Wesleyan University*

Tian-Xiao He, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

---

Nathan Mueggenburg and Tian-Xiao He, Faculty Advisor, "Adapting Daubechies' Wavelet Iteration Scheme to Biorthogonal Scaling Functions" (April 12, 1997). *John Wesley Powell Student Research Conference*. Paper 4. <http://digitalcommons.iwu.edu/jwprc/1997/oralpres2/4>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

## Oral Presentation 2.2

### ADAPTING DAUBECHIES' WAVELET ITERATION SCHEME TO BIORTHOGONAL SCALING FUNCTIONS

Nathan Mueggenburg and Tian-Xiao He\*, Department of Mathematics, IWU

In 1988 Daubechies developed an iteration scheme to produce a D4 wavelet function which has since become very popular. This iteration scheme is based upon four conditions stated by Daubechies. One of these conditions required that the scaling function be orthogonal. We propose a new iteration scheme, similar in nature to Daubechies', but with a relaxed orthogonality condition. Our iteration scheme does not require that the scaling function be orthogonal, but only that it be biorthogonal. With this orthogonality condition relaxed, we can construct a dual of an unorthogonal function by using macroelement analysis, and thus create a larger set of possible initial functions for the iteration scheme. This new iteration scheme is then able to produce biorthogonal wavelet functions, a much broader group than that produced by Daubechies' iteration scheme. As an example we use the second order B-spline, commonly known as the hat function, as an initial function of our iteration scheme.